

---

csci54 – discrete math & functional programming  
propositional logic

---

---


Simplify each of the following Haskell expressions:

(a) `a && not a`

(b) `a || (not a && b)`


(c) `(not a || b) && (not b || c) &&  
     (not c || not a) && (not c || not b)`





George  
Boole  
1815-1864

---



# On "True" and "False"

---

- ▶ logic is the study of valid reasoning
  
- ▶ The starting point:  
    A proposition is a statement that is either True or False.
  
- ▶ What are examples of propositions that are True? False? Unknown?



# On propositional logic

---

- ▶ the study of propositions: how to formulate, evaluate, manipulate
- ▶ atomic proposition: a proposition that is conceptually indivisible
- ▶ compound proposition: a proposition that is build up out of conceptually simpler propositions
  - ▶ How?



# Creating compound propositions

---

- ▶ We can create more complex propositional statements using logical connectives

- ▶ negation (not,  $\neg$ ,  $\sim$ )
- ▶ conjunction (and,  $\wedge$ )
- ▶ disjunction (or,  $\vee$ )
- ▶ implication (implies,  $\Rightarrow$ ,  $\rightarrow$ )

## Precedence rules:

- negation binds most tightly
- then conjunction
- then disjunction
- then implication

implication is right-associative

- ▶ In particular, a well-formed propositional logic formula is defined as:

$$\phi ::= T | F | (\neg\phi) | (\phi \wedge \phi) | (\phi \vee \phi) | (\phi \Rightarrow \phi)$$

---



# Evaluating compound propositional statements

- ▶ Convenient to use a truth table to display the relationships between truth values of different propositions

- ▶ Truth table for negation: 

$p$	$\neg p$
$T$	$F$
$F$	$T$

- ▶ For conjunction (and) and disjunction (or): 

$p$	$q$	$p \wedge q$	$p \vee q$
$T$	$T$	$T$	$T$
$T$	$F$	$F$	$T$
$F$	$T$	$F$	$T$
$F$	$F$	$F$	$F$

$$\phi ::= T | F | (\neg \phi) | (\phi \wedge \phi) | (\phi \vee \phi) | (\phi \Rightarrow \phi)$$



# Implication

---

- ▶ What does it mean to say "p implies q"?

- ▶ p  $\Rightarrow$  q is true if q is true or p is false

p	q	p $\Rightarrow$ q
T	T	T
T	F	F
F	T	T
F	F	T

- ▶ What is the truth value of each of the following statements?

- ▶  $1 + 1 = 2$  implies that  $2 + 3 = 5$
- ▶  $1 + 1 = 2$  implies that  $2 + 3 = 6$
- ▶  $1 + 1 = 3$  implies that  $2 + 3 = 5$
- ▶  $1 + 1 = 3$  implies that  $2 + 3 = 6$





# A little more on implications

---

▶  $p \Rightarrow q$

- ▶ “if p, then q”
- ▶ “p implies q”
- ▶ “p only if q”
- ▶ “q whenever p”
- ▶ “q, if p”
- ▶ “q is necessary for p”
- ▶ “p is sufficient for q”

▶ Bidirectional implication  $p \Leftrightarrow q$

- ▶ “p if and only if q”, “p iff q”
- ▶ True only when p and q have same truth value: either both true or both false.

# Example

---

- ▶ "Since Sandra is wearing a soccer jersey, she must be a soccer player."
- ▶ This compound proposition is composed of 2 atomic propositions:
  - ▶ (1) = Sandra is wearing a soccer jersey
  - ▶ (2) = Sandra is a soccer player
- ▶ The compound proposition can be written as:
  - ▶ (1)  $\leftrightarrow$  (2)

# Passwords

---

- ▶ "A password is valid only if it is at least 8 characters long, is not one that you have used previously, and contains at least 2 of the following: a number, a lowercase character, an uppercase character."
- ▶ This is a compound proposition that is composed of how many atomic propositions?
- ▶ What are the 6 atomic propositions?
- ▶ How can you write the compound proposition in terms of the atomic propositions?



---

---



## categorizing well-formed formulas (wff)

---

- ▶ A formula in propositional logic is one of:
  - ▶ tautology (valid): if it evaluates to T in all cases
  - ▶ satisfiable: evaluates to T in some cases
  - ▶ contingency (falsifiable): evaluates to F in some cases
  - ▶ contradiction (unsatisfiable): evaluates to F in all cases

- ▶ Consider the following formula:

$$(p \vee q) \Rightarrow (\neg p \wedge \neg q)$$

- ▶ Which of the following describes the formula: tautology, satisfiable, contingency, contradiction? Why?



# a collection of tautologies

---

---

$(p \Rightarrow q) \wedge p \Rightarrow q$       Modus Ponens

$(p \Rightarrow q) \wedge \neg q \Rightarrow \neg p$       Modus Tollens

---

$p \vee \neg p$       Law of the Excluded Middle

$p \Leftrightarrow \neg\neg p$       Double Negation

$p \Leftrightarrow p$

---

$p \Rightarrow p \vee q$

$p \wedge q \Rightarrow p$

---

---

$(p \vee q) \wedge \neg p \Rightarrow q$

$(p \Rightarrow q) \wedge (\neg p \Rightarrow q) \Rightarrow q$

$(p \Rightarrow q) \wedge (q \Rightarrow r) \Rightarrow (p \Rightarrow r)$

---

$(p \Rightarrow q) \wedge (p \Rightarrow r) \Leftrightarrow p \Rightarrow q \wedge r$

$(p \Rightarrow q) \vee (p \Rightarrow r) \Leftrightarrow p \Rightarrow q \vee r$

$p \wedge (q \vee r) \Leftrightarrow (p \wedge q) \vee (p \wedge r)$

$p \Rightarrow (q \Rightarrow r) \Leftrightarrow p \wedge q \Rightarrow r$

---



# logical equivalence

---

- ▶ Two propositions are logically equivalent ( written ) if they have exactly identical truth tables (i.e. their truth values are the same under every truth assignment)

Simplify each of the following Haskell expressions:

(a) `a && not a`

(b) `a || (not a && b)`

(c) `(not a || b) && (not b || c) &&  
(not c || not a) && (not c || not b)`



# some logically equivalent propositions

---

Commutativity

$$p \vee q \equiv q \vee p$$

$$p \wedge q \equiv q \wedge p$$

$$p \oplus q \equiv q \oplus p$$

$$p \Leftrightarrow q \equiv q \Leftrightarrow p$$

---

Associativity

$$p \vee (q \vee r) \equiv (p \vee q) \vee r$$

$$p \wedge (q \wedge r) \equiv (p \wedge q) \wedge r$$

$$p \oplus (q \oplus r) \equiv (p \oplus q) \oplus r$$

$$p \Leftrightarrow (q \Leftrightarrow r) \equiv (p \Leftrightarrow q) \Leftrightarrow r$$

---

Idempotence

$$p \vee p \equiv p$$

$$p \wedge p \equiv p$$

Distribution of  $\wedge$  over  $\vee$

$$p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$$

Distribution of  $\vee$  over  $\wedge$

$$p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$$

---

Contrapositive

$$p \Rightarrow q \equiv \neg q \Rightarrow \neg p$$

---

$$p \Rightarrow q \equiv \neg p \vee q$$

$$p \Rightarrow (q \Rightarrow r) \equiv p \wedge q \Rightarrow r$$

$$p \Leftrightarrow q \equiv \neg p \Leftrightarrow \neg q$$

---

Mutual Implication  $(p \Rightarrow q) \wedge (q \Rightarrow p) \equiv p \Leftrightarrow q$

---

De Morgan's Laws

$$\neg(p \wedge q) \equiv \neg p \vee \neg q$$

$$\neg(p \vee q) \equiv \neg p \wedge \neg q$$

$$(\neg a \vee b) \wedge (\neg b \vee c) \wedge (\neg c \vee \neg a) \wedge (\neg c \vee \neg b)$$

---

