
csci54 – discrete math & functional programming
propositional logic continued, predicate logic

last time

- ▶ introduction to propositional logic:

- ▶ Boole

- ▶ proposition

- ▶ well-formed propositional logic formulas (wff)

$$\phi ::= T|F|(\neg\phi)|(\phi \wedge \phi)|(\phi \vee \phi)|(\phi \Rightarrow \phi)$$

- ▶ truth tables for operators

- ▶ tautology/satisfiable/contingency (falsifiable)/contradiction

- ▶ implication

- ▶ logical equivalence



converse, inverse, contrapositive

Given an implication $p \Rightarrow q$, we can derive three other implications:

- converse: $q \rightarrow p$
 - inverse: $\neg p \rightarrow \neg q$
 - contrapositive: $\neg q \rightarrow \neg p$
- Which, if any, of the converse, inverse, and contrapositive is logically equivalent to the original implication?





consider the following statements . . .

- ▶ If 2 is an even number then 3 is an odd number.
- ▶ If x is an even number, then $x+1$ is an odd number.

- ▶ How would you express these two statements in propositional logic?



predicate logic

- ▶ A predicate P is function that assigns the value True or False to each element of a set U .
 - ▶ The set U is called the universe or domain of discourse
 - ▶ P is a predicate over U
- ▶ Examples:
 - ▶ the predicate "is an even number" over the positive integers.
 - ▶ the predicate "last name has at least 6 characters" over the set of people currently in this room.
- ▶ Once you specify the element of U , then you have a proposition with a truth value.



quantifiers

- ▶ quantifiers are another way to form propositions from a predicate

Definition 3.21: Universal quantifier [for all, \forall].

Let P be a predicate over S . The proposition $\forall x \in S : P(x)$ is true if, for *every* possible $x \in S$, we have that $P(x)$ is true.

Definition 3.22: Existential quantifier [there exists, \exists].

Let P be a predicate over S . The proposition $\exists x \in S : P(x)$ is true if, for *at least one* possible $x \in S$, we have that $P(x)$ is true.



quantifiers - example

- ▶ Imagine these predicates
 - ▶ "rested(n)" means "n got at least 8 hours of sleep in the past 24 hours"
 - ▶ "bornMA(n)" means "n was born in Massachusetts"
- ▶ Which, if any, of the following propositions is true? Justify your answer.
 - ▶ $\forall n$ in this room : rested(n)
 - ▶ $\forall n$ in this room : (rested(n) \wedge bornMA(n))
 - ▶ $\exists n$ currently enrolled at Pomona College : (rested(n) \vee bornMA(n))
 - ▶ $\exists n$ currently enrolled at Pomona College : (rested(n) \wedge bornMA(n))



free and bound variables (an aside)

- ▶ In an expression variables can be free/unbound or bound
 - ▶ With a free variable the value is not fixed by the expression
 - ▶ With a bound variable the value is defined within the expression

$$\forall x \in \mathbb{Z} : x^2 \geq y$$

- ▶ An expression of predicate logic with no free variables is called fully quantified



theorems in predicate logic

- ▶ A fully quantified expression of predicate logic is a theorem if and only if it is true for every possible meaning of each of its predicates.
- ▶ Is the following a theorem?
$$[\forall x \in S : P(x)] \vee [\forall x \in S : \neg P(x)]$$
- ▶ What is an example of a predicate for which the statement is false? is true?





practice question

- ▶ Exactly one of the following two propositions is a theorem.
Which one?

$$(1) \quad [\forall x \in S : P(x) \vee Q(x)] \Leftrightarrow [\forall x \in S : P(x)] \vee [\forall x \in S : Q(x)]$$

$$(2) \quad [\exists x \in S : P(x) \vee Q(x)] \Leftrightarrow [\exists x \in S : P(x)] \vee [\exists x \in S : Q(x)]$$

- ▶ Justify your answer